

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau






(43) International Publication Date
15 February 2001 (15.02.2001)

PCT

(10) International Publication Number
WO 01/10882 A1

- (51) International Patent Classification⁷: C07H 19/10, 19/20
- (21) International Application Number: PCT/CA00/00914
- (22) International Filing Date: 10 August 2000 (10.08.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/147,820 10 August 1999 (10.08.1999) US
- (71) Applicant (for all designated States except US): ADANI, Alexander [CA/CA]; 126 Pine Valley Crescent, Woodbridge, Ontario L4L 2W4 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): SHIROKOVA, Elena A. [RU/RU]; Akademika Artsymovicha, str. 9, app. 17, Moscow, 117437 (RU). KHANDAZHYNSKAYA, Anastasiya L. [RU/RU]; 3-Frunzenskaya, str. 12, app. 27, Moscow, 119146 (RU). KARPENKO, Inna L. [RU/RU]; Shebalinsky proezd, 8, app. 11, Moscow, 108125 (RU). TARUSOVA, Natalya B. [RU/RU]; Zhivopisnaya, str. 50, app. 40, Moscow, 123098 (RU). KRAYEVSKY, Alexander A. [RU/RU]; Profsoyuznaya str. 132/4, app. 111, Moscow, 117321 (RU).
- (74) Agents: NASSIF, Omar, A. et al.; Gowling, Strathy & Henderson, Suite 4900, Commerce Court West, Toronto, Ontario M5L 1J3 (CA).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— With international search report.
— Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DEPOT FORMS OF MODIFIED NUCLEOSIDE 5'-HYDROGENPHOSPHATES AS INHIBITORS OF REPRODUCTION OF HUMAN IMMUNODEFICIENCY VIRUS AND HUMAN HEPATITIS B VIRUS



(57) Abstract: The present invention is directed to novel compounds exhibiting a selective inhibition of the reproduction of HIV and HBV and possessing low toxicity. The present compounds are VI-VIII of formulae (VI, VII and VIII) wherein: Nucleosides of D and L-series; B = thymine, adenine, guanine, cytosine, uracyl, 5-fluorouracyl or 5-ethyluracyl; R = N₃, NH₂, F, H or F; X = S or O; R = adamantyl-1, bicyclo[6,5,1] heptyl-1 or *tert*-butyl. These compounds being depot forms of 5'-hydrogenphosphonates of modified nucleosides are able to inhibit the reproduction of HIV and HBV and are less toxic as compared to the prior art compounds.

WO 01/10882 A1

DEPOT FORMS OF MODIFIED NUCLEOSIDE 5'-HYDROGENPHOSPHATES AS INHIBITORS OF REPRODUCTION OF HUMAN IMMUNODEFICIENCY VIRUS AND HUMAN HEPATITIS B VIRUS

FIELD OF THE INVENTION

The present invention relates to the art of molecular biology and virology. More specifically, in one of its aspects, the present invention relates to novel depot forms of modified 5'-hydrogenphosphonates as selective inhibitors of the reproduction of Human Immunodeficiency Virus (HIV) and Human Hepatitis B Virus (HBV) in cell cultures. In another of its aspects, the present invention relates to a process for producing depot forms of modified 5'-hydrogenphosphonates.

10 DESCRIPTION OF THE PRIOR ART

It is known in the art that various 5'-hydrogenphosphonates of 3'-azido-3'-deoxythymidine (phosphazide) [1-3] and other modified nucleosides [4-6] inhibit the reproduction of the HIV. While somewhat effective, these compounds are comparatively hydrophilic because of the presence of the unprotected phosphonate group thereby decreasing diffusion thereof into cells.

To the knowledge of the present inventors, only one attempt has been made to prepare a depot form of phosphazide [7], but in human serum these compounds are hydrolyzed to 3'-azido-3'-deoxythymidine rather than to phosphazide. This pathway dominates over the other due to hydrolysis of the nucleoside-phosphonate bond proceeding more rapidly than the elimination of the phosphonate protective group. An example of a depot form of 2',3'-dideoxy-2',3'-didehydrothymidine hydrogenphosphonate has been published [8], but the structure of this compound was not taught.

25 SUMMARY OF THE INVENTION

We have discovered a principal for the design of depot forms of nucleoside 5'-hydrogenphosphonates and a group of depot forms of modified nucleoside 5'-hydrogenphosphonates having anti HIV and HBV activity. The compounds according to the present invention, viz. depot forms of modified nucleoside 5'-hydrogenphosphonates, are able to inhibit selectively the reproduction of HIV and HBV in cell cultures.

DEPOT FORMS OF MODIFIED NUCLEOSIDE 5'-HYDROGENPHOSPHATES AS INHIBITORS OF REPRODUCTION OF HUMAN IMMUNODEFICIENCY VIRUS AND HUMAN HEPATITIS B VIRUS

FIELD OF THE INVENTION

The present invention relates to the art of molecular biology and virology. More specifically, in one of its aspects, the present invention relates to novel depot forms of modified 5'-hydrogenphosphonates as selective inhibitors of the reproduction of Human Immunodeficiency Virus (HIV) and Human Hepatitis B Virus (HBV) in cell cultures. In another of its aspects, the present invention relates to a process for producing depot forms of modified 5'-hydrogenphosphonates.

10 DESCRIPTION OF THE PRIOR ART

It is known in the art that various 5'-hydrogenphosphonates of 3'-azido-3'-deoxythymidine (phosphazide) [1-3] and other modified nucleosides [4-6] inhibit the reproduction of the HIV. While somewhat effective, these compounds are comparatively hydrophilic because of the presence of the unprotected phosphonate group thereby decreasing diffusion thereof into cells.

To the knowledge of the present inventors, only one attempt has been made to prepare a depot form of phosphazide [7], but in human serum these compounds are hydrolyzed to 3'-azido-3'-deoxythymidine rather than to phosphazide. This pathway dominates over the other due to hydrolysis of the nucleoside-phosphonate bond proceeding more rapidly than the elimination of the phosphonate protective group. An example of a depot form of 2',3'-dideoxy-2',3'-didehydrothymidine hydrogenphosphonate has been published [8], but the structure of this compound was not taught.

25 SUMMARY OF THE INVENTION

We have discovered a principal for the design of depot forms of nucleoside 5'-hydrogenphosphonates and a group of depot forms of modified nucleoside 5'-hydrogenphosphonates having anti HIV and HBV activity. The compounds according to the present invention, viz. depot forms of modified nucleoside 5'-hydrogenphosphonates, are able to inhibit selectively the reproduction of HIV and HBV in cell cultures.


THIS PAGE BLANK (USPTO)

According to the synthetic pathway set out in Figure 1, prodrug I can be hydrolyzed in two possible ways: to nucleoside 5'-hydrogenphosphonate (II, Route a) or nucleoside III (Route b). The route of choice, *inter alia*, depends on the stability and steric availability of the corresponding bonds in I. Our study of several groups of prodrugs in human serum showed that most compounds were hydrolyzed via Route b except for those bearing tertiary alkyl esters, which were hydrolyzed via Route a.

The third possibility is the preliminary oxidation of I to IV via Route c with subsequent hydrolysis either to V (via Route d) or to III (via Route e). It is known that oxidation proceeds much faster for hydrogenphosphonate diesters than for hydrogenphosphonate monoesters.

The study of diesters of nucleoside hydrogenphosphonates in human serum showed that most compounds were hydrolyzed via Route b except for those bearing tertiary alkyl esters, which were hydrolyzed via Route a.

Prodrug forms of 5'-hydrogenphosphonates of modified nucleosides hydrolyzed to the corresponding nucleoside 5'-hydrogenphosphonates include:



wherein:

nucleosides of D and L-series;

B = thymine, adenine, guanine, cytosine, uracil, 5-fluorouracil or 5-ethyluracil;

R = N₃, H or F;

X = S or O; and

R' = adamantyl-1, bicyclo[2,2,1]heptyl-1, *tert.*-butyl.


BRIEF DESCRIPTION OF THE DRAWINGS

Embodiments of the present invention will be described with reference to the accompanying drawing, in which:

Figure 1 illustrates a synthetic pathway, inter alia, for the production of prodrug I.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The synthesis of VI-VIII with thymine and adenine bases may be achieved according to the following reaction scheme (referred to hereinbelow as Scheme 2):



wherein:

NUCLEOSIDE = nucleoside component;

R' = adamantyl-1, bicyclo[2,2,1] heptyl-1 or *tert.*-butyl; and

R'' = COC(CH₃)₃, -SO₂CH₃ or -SO₂CF₃.

The synthesis of VI-VIII with thymine, adenine, and guanine bases may be achieved according to the following reaction scheme (referred to hereinbelow as Scheme

3):

R' = adamantyl-1, bicyclo[2,2,1]heptyl-1, *tert.*-butyl.

BRIEF DESCRIPTION OF THE DRAWINGS


Embodiments of the present invention will be described with reference to the accompanying drawing, in which:

Figure 1 illustrates a synthetic pathway, inter alia, for the production of prodrug

I.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The synthesis of VI-VIII with thymine and adenine bases may be achieved according to the following reaction scheme (referred to hereinbelow as Scheme 2):



wherein:

NUCLEOSIDE = nucleoside component;


R' = adamantyl-1, bicyclo[2,2,1] heptyl-1 or *tert.*-butyl; and

R'' = COC(CH₃)₃, -SO₂CH₃ or -SO₂CF₃.

The synthesis of VI-VIII with thymine, adenine, and guanine bases may be achieved according to the following reaction scheme (referred to hereinbelow as Scheme

3):

THIS PAGE BLANK (USPTO)




wherein:

NUCLEOSIDE = nucleoside component; and

R' = adamantyl-1, bicyclo[2,2,1]heptyl-1 or *tert.*-butyl.

The synthesis of **VI-VIII** with different nucleic bases may be achieved according the following reaction scheme (referred to hereinbelow as Scheme 4):



wherein:

NUCLEOSIDE = nucleoside component;


R' = adamantyl-1, bicyclo[2,2,1] heptyl-1 or *tert.*-butyl.


5 Embodiments of the present invention will be described with reference to the following Examples which are provided for illustrative purposes only and should not be use to construe or limited the scope of the present invention.

[A] EXAMPLES OF THE SYNTHETIC PROCEDURE

10

EXAMPLE 1: P-(Adamantyl-1) 3'-azido-2',3'-dideoxythymidine 5'-hydro-
genphosphonate (VIa, B=Thy, R=N₃, R'=adamantyl-1) - Scheme 2





wherein:

NUCLEOSIDE = nucleoside component;


R' = adamantyl-1, bicyclo[2,2,1] heptyl-1 or *tert.*-butyl.

5 Embodiments of the present invention will be described with reference to the following Examples which are provided for illustrative purposes only and should not be use to construe or limited the scope of the present invention.

[A] EXAMPLES OF THE SYNTHETIC PROCEDURE

10

EXAMPLE 1: P-(Adamantyl-1) 3'-azido-2',3'-dideoxythymidine 5'-hydro-
genphosphonate (VIa, B=Thy, R=N₃, R'=adamantyl-1) - Scheme 2




THIS PAGE BLANK (USPTO)

A water solution (0.5 ml) of 3'-azido-2',3'-dideoxythymidine 5'-hydrogenphosphonate (Phosphazide) Na-salt (88 mg, 0.25 mmol) was passed through a Dowex-50 (Py⁺) column (5 x 1 cm), the column washed with water until no UV-adsorption at 260 nm was observed. The eluate was evaporated to dryness and coevaporated with pyridine (3 x 2 ml). The residue was dissolved in MeCN (5 ml), cooled to -20°C and adamantanol-1 (58 mg, 0.4 mmol) was added under stirring. Pyridine (0.5 ml) and pivaloyl chloride (90 mg, 91 ml, 0.75 mmol) were added, cooling was removed and the reaction mixture was stirred for 10 min. The mixture was diluted with chloroform (10 ml) and washed with cooled saturated sodium bicarbonate (5 ml) and water (3 x 3 ml). The organic solution was dried with Na₂SO₄, evaporated, and reevaporated with toluene. The product was purified on a Kieselgel 60 column (15 x 2.5 cm) eluting with chloroform : methanol 95 : 5. The target fractions were evaporated to give compound VIa. The yield and physicochemical data are given in Tables 1-3.

Example 2: P-(Adamantyl-1) 3'-azido-2',3'-dideoxythymidine 5'-hydrogenphosphonate (VIa, B=Thy, R=N₃, R'=adamantyl-1) - Scheme 3

A solution of PCl₃ (75 mg, 44 ml, 0.5 mmol) in dichloromethane (2 ml) was cooled to 1-2°C, adamantanol-1 (79 mg, 0.5 mmol) and pyridine (40 ml, 0.5 mmol) were added for 20 min at 4-5°C under stirring. A solution of 3'-azido-2',3'-dideoxythymidine prepared from Na-salt (65 mg, 0.25 mmol) as in the example 1 in pyridine (2 ml) was dropped for 3 h. The reaction mixture was stirred for 3 h at 18°C and then quenched with chloroform (5 ml) and a cooled solution of 1.5 M NH₄HCO₃ (3 ml), the organic layer was washed with water, dried with sodium sulfate, and evaporated. The residue was purified on a Kieselgel 60 column (2 x 25 mm) eluting with CHCl₃-tert.-BuOH 9 : 1 to give compound VIa. The yield and physicochemical data are given in Tables 1-3.


Example 3: P-tert.-Butyl 3'-azido-2',3'-dideoxythymidine 5'-hydrogenphosphonate (VIb, B=Thy, R=N₃, R'=tert.-butyl) - Scheme 2



Phosphazide Na-salt (88 mg, 0.25 mmol) was coupled with *tert.*-butanol (24 mg, 35 ml, 0.38 mmol) under the conditions of example 1. For the yield and physicochemical data, see Tables 1-3.

5

Example 4: *P*-(Adamantyl-1)2',3'-dideoxythymidine 5'-hydrogenphosphonate (VIc, B=Thy, R=H, R'=adamantyl-1) - Scheme 2.




10

2',3'-Dideoxythymidine 5'-hydrogenphosphonate Na-salt (78 mg, 0.25 mmol) was coupled with adamantanol-1 (58 mg, 0.4 mmol) under the conditions of Example 1 to give VIc. For the yield and physicochemical data, see Tables 1-3.

15 Example 5: *P*-(Adamantyl-1)2',3'-dideoxythymidine 5'-hydrogenphosphonate (VIc, B=Thy, R=H, R'=adamantyl-1) - Scheme 4.


20 A solution of 2',3'-dideoxythymidine 5'-hydrogenphosphonate (65 mg, 0.25 mmol) in dichloromethane (1 ml) and pyridine (1 ml) was added to the solution of diphenylphosphite (97 mg, 0.5 mmol) in acetonitrile (2 ml) at 0°C and, after 30-min stirring, adamantanol-1 (58 mg, 0.4 mmol) was added to the reaction mixture under stirring. After 1 h, the mixture was diluted with triethylamine - water 1 : 1, evaporated,



Phosphazide Na-salt (88 mg, 0.25 mmol) was coupled with *tert.*-butanol (24 mg, 35 ml, 0.38 mmol) under the conditions of example 1. For the yield and physicochemical data, see Tables 1-3.

5

Example 4: *P*-(Adamantyl-1)2',3'-dideoxythymidine 5'-hydrogenphosphonate (VIc, B=Thy, R=H, R'=adamantyl-1) - Scheme 2.



10

2',3'-Dideoxythymidine 5'-hydrogenphosphonate Na-salt (78 mg, 0.25 mmol) was coupled with adamantanol-1 (58 mg, 0.4 mmol) under the conditions of Example 1 to give VIc. For the yield and physicochemical data, see Tables 1-3.

15 **Example 5:** *P*-(Adamantyl-1)2',3'-dideoxythymidine 5'-hydrogenphosphonate (VIc, B=Thy, R=H, R'=adamantyl-1) - Scheme 4.


20 A solution of 2',3'-dideoxythymidine 5'-hydrogenphosphonate (65 mg, 0.25 mmol) in dichloromethane (1 ml) and pyridine (1 ml) was added to the solution of diphenylphosphite (97 mg, 0.5 mmol) in acetonitrile (2 ml) at 0°C and, after 30-min stirring, adamantanol-1 (58 mg, 0.4 mmol) was added to the reaction mixture under stirring. After 1 h, the mixture was diluted with triethylamine - water 1 : 1, evaporated,

THIS PAGE BLANK (USPTO)

the residue was diluted with chloroform (3 ml), washed with water (3 x 1 ml), and the residue chromatographed on a Kieselgel 60 column (2 x 25 mm) eluting with CHCl₃-ethanol 9 : 1 to give compound VIc. The yield and physicochemical data are given in Tables 1-3.

5


Example 6: P-(Adamantyl-1) 2',3'-dideoxycytidine 5'-hydrogenphosphonate (VIId, B=Cyt, R=H, R'=adamantyl-1). Scheme 4



10 2',3'-Dideoxycytidine (105 mg, 0.5 mmol), adamantanol-1 (116 mg, 0.8 mmol), and diphenylphosphite (194 mg, 1 mmol) were coupled under the conditions of example 5. For the yield and physicochemical data of VIId, see Tables 1-3.

15


Example 7: P-(Adamantyl-1) 3'-fluoro-2',3'-dideoxythymidine 5'-hydrogenphosphonate (VIe, B=Thy, R=F, R'=adamantyl-1) - Scheme 2



20


3'-Fluoro-2',3'-dideoxythymidine 5'-hydrogenphosphonate Na-salt (82 mg, 0.25 mmol) was converted to VIe as described in example 1. The yield and physicochemical data are given in Tables 1-3.

Example 8: P-(Adamantyl-1) 2',3'-dideoxyadenosine 5'-hydrogenphosphonate (VIIf, B=Ade, R=H, R'=Adamantyl-1) - Scheme 4



2',3'-Dideoxyadenosine (118 mg, 0.5 mmol) was converted to **VIg** under the conditions of example 6. The yield and physicochemical data of **VIg** are given in Tables 1-3.


- 5 **Example 9:** *P-tert.-Butyl 2',3'-dideoxyadenosine 5'-hydrogenphosphonate (VIg, B=Ade, R=H, R'=tert.-butyl)* - Scheme 4.



2',3'-Dideoxyadenosine (118 mg, 0.5 mmol) was converted to **VIg** under the conditions of example 6. The yield and physicochemical data are given in Tables 1-3.

10


- Example 10:** *P-(Adamantyl-1) 2',3'-dideoxyguanosine 5'-hydrogenphosphonate (VIh, B=Gua, R=H, R'=adamantyl-1)* - Scheme 4



- 15 2',3'-Dideoxyguanosine (125 mg, 0.5 mmol) was converted to **VIh**, as in the **Example 6**. The yield and physicochemical data are given in Tables 1-3.


- Example 11:** *P-(Adamantyl-1) 2',3'-dideoxy-2',3'-didehydrothymidine 5'-hydrogenphosphonate (VIIa, B=Thy, R'=adamantyl-1)* - Scheme 2

20



2',3'-Dideoxyadenosine (118 mg, 0.5 mmol) was converted to **VI f** under the conditions of example 6. The yield and physicochemical data of **VI f** are given in Tables 1-3.


- 5 **Example 9:** *P-tert.-Butyl 2',3'-dideoxyadenosine 5'-hydrogenphosphonate (VI g, B=Ade, R=H, R'=tert.-butyl)* - Scheme 4.



2',3'-Dideoxyadenosine (118 mg, 0.5 mmol) was converted to **VI g** under the conditions of example 6. The yield and physicochemical data are given in Tables 1-3.

10

- Example 10:** *P-(Adamantyl-1) 2',3'-dideoxyguanosine 5'-hydrogenphosphonate (VI h, B=Gua, R=H, R'=adamantyl-1)* - Scheme 4




- 15 2',3'-Dideoxyguanosine (125 mg, 0.5 mmol) was converted to **VI h**, as in the Example 6. The yield and physicochemical data are given in Tables 1-3.

- Example 11:** *P-(Adamantyl-1) 2',3'-dideoxy-2',3'-didehydrothymidine 5'-hydrogenphosphonate (VII a, B=Thy, R'=adamantyl-1)* - Scheme 2

20


THIS PAGE BLANK (USPTO)



2',3'-Dideoxy-2',3'-didehydrothymidine 5'-hydrogenphosphonate Na-salt (78 mg, 0.25 mmol) was converted to **VIIa** under the conditions of example 1. The yield and physicochemical data are given in Tables 1-3.

5


Example 12: P-(Adamantyl-1)-2',3'-dideoxy-2',3'-didehydrocytidine 5'-hydrogenphosphonate (VIIb B=Cyt, R'=adamantyl-1) - Scheme 4



10


VIIb was synthesized from 2',3'-dideoxy-2',3'-didehydrocytidine (125 mg, 0.5 mmol) under the conditions of Example 6. The yield and physicochemical data are given in Tables 1-3.

15 *Example 13:* P-(Adamantyl-1)-L-3'-thiocytidine 5'-hydrogenphosphonate (VIIIa, B=Cyt, R'=adamantyl-1) - Scheme 3



Compound VIIIa was synthesized from L-3'-thiocytidine (120 mg, 0.5 mmol) under the conditions of example 3. The yield and physicochemical data are given in Tables 1-3.

- 5 **Example 14:** *P-tert.-Butyl L-3'-thiocytidine 5'-hydrogenphosphonate (VIIIb, B=Cyt, R'=tert.-butyl)* - Scheme 4



- 10 Compound VIIIb was synthesized from L-3'-thiocytidine (120 mg, 0.5 mmol) according to Example 3. The yield and physicochemical data are given in Tables 1-3.


TABLE 1

Yields and physicochemical characteristics

Compound	R	Mol. mass	Yield, %	UV, MeOH, λ , nm
VIa	adamantyl-1	463	67	264
VIb	tert.-butyl	387	76	265
VIc	adamantyl-1	421	78	264
VI d	adamantyl-1	407	61	274
VIe	adamantyl-1	439	88	265
VI f	adamantyl-1	431	65	257
VI g	tert.-butyl	355	59	257
VI h	adamantyl-1	447	59	250
VIIa	adamantyl-1	419	79	265
VIIb	adamantyl-1	415	62	274
VIIIa	adamantyl-1	425	57	272
VIIIb	tert.-butyl	349	48	273

Compound VIIIa was synthesized from L-3'-thiocytidine (120 mg, 0.5 mmol) under the conditions of example 3. The yield and physicochemical data are given in Tables 1-3.

- 5 *Example 14: P-tert.-Butyl L-3'-thiocytidine 5'-hydrogenphosphonate (VIIIb, B=Cyt, R'=tert.-butyl) - Scheme 4*



- 10 Compound VIIIb was synthesized from L-3'-thiocytidine (120 mg, 0.5 mmol) according to Example 3. The yield and physicochemical data are given in Tables 1-3.

TABLE 1

Yields and physicochemical characteristics

Compound	R	Mol. mass	Yield, %	UV, MeOH, λ_{max} , nm
VIa	adamantyl-1	463	67	264
VIb	tert.-butyl	387	76	265
VIc	adamantyl-1	421	78	264
VIId	adamantyl-1	407	61	274
VIe	adamantyl-1	439	88	265
VIIf	adamantyl-1	431	65	257
VIg	tert.-butyl	355	59	257
VIh	adamantyl-1	447	59	250
VIIa	adamantyl-1	419	79	265
VIIb	adamantyl-1	415	62	274
VIIIa	adamantyl-1	425	57	272
VIIIb	tert.-butyl	349	48	273

THIS PAGE BLANK (USPTO)

TABLE 2.

¹H NMR spectra of VI-VIII, 200 MHz; CD₃CN; δ, ppm; (J, Hz):

Comp.	H1' 1H	H2'	H3' 1H	H4' 1H	H5' 2H	H-5 (pyr) or H-2 (pur) 1H	H-6 (pyr) or H-8 (pur) 1H	CH ₃ , Thy	P ³¹
VIa	6.14dd (7)	2.38m 2H	4.34m	4.01m	4.15m	-	7.38 q (1)	1.96d (7)	6.93d (696)
VIb	6.15dd (6.5)	2.38m, 2H	4.37m	4.01m	4.20m	-	7.38q (1)	1.8d (6.5)	6.90d (697)
VIc	6.05m	4.4m, 2H	5.03t (10.7)	4.2m	3.62q (6.55)	-	6.45d (7.3)	1.7d (7.7)	6.48d (638)
VI d	6.4m	2.15m, 2H	3.52d (10.7)	4.2m	3.99m	6.35d (7.7)	8.04d (7.7)	-	6.71d (630)
VIe	6.21t (6.6)	2.46t (6.2), 2H	4.63t (5.43)	3.38s	4.2m	-	7.65d (0.5)	1.8d (0.8)	7.11d (638)
VI f	6.34q (7)	2.14m, 2H	2.66m	4.24m	3.56m	8.39s	8.16s	-	6.59d (638)
VI g	6.28q (6.5)	2.26t (6.4), 2H	2.76m	4.28m	3.48m	8.46s	8.04s	-	6.66d (641)
VI h	6.04q (6)	2.12m, 2H	2.49m	4.1m	3.9m	8.00c	-	-	6.56d (632)
VIIa	6.95t (5)	4.98m, 1H	5.25m	4.05m	3.65m	6.45d (7.3)	7.72d (7.3)	-	6.48d (638)
VIIb	6.95t (5.2)	5.12m, 1H	5.31m	4.11m	3.55m	6.35d (7.6)	8.04d (7.7)	-	6.71d (630)
VIIIa	6.6dd (3.0,5.9)	3.4dd (3.0; 5.7), 2H	-	5.24t (4.6)	3.65m 2H	5.88d (7.7)	7.89d (7.7)	-	6.71d (630)
VIIIb	6.52dd (3.1,5.8)	3.48dd (3.0; 5.6), 2H	-	5.24t (4.6)	3.72m 2H	5.80d (7.7)	7.75d (7.7)	-	6.71d (630)

TABLE 3

¹H NMR spectral data for "depot" groups in VI-VIII, 200 MHz; CD₃CN; δ, ppm; (J, Hz) and
³¹P NMR spectra of VI-VIII, 81 MHz, CD₃CN; δ, ppm, (J, Hz)

Comp.	<i>tert.</i> -CH (adam.), 3H	CH ₂ -C-O- P(adam.), 6H	CH ₂ (adam.) 6H	<i>tert.</i> -Bu, 9H	³¹ P NMR, 1P
VIa	2.16 br.s	2.06m	1.64m	-	3.36d (696), 3.08d
VIb	-	-	-	1.49s	3.99dt (697 and 8), 3.63 dt.
VIc	2.16 br.s	2.06m	1.63m	-	3.38d (694), 3.04d
VI d	2.16 br.s	2.08m	1.64m	-	3.33d (696), 3.07d
VIe	2.16 br.s	2.08m	1.64m	-	3.36d (694), 3.06d
VI f	2.16 br.s	2.06m	1.63m	-	3.32d (695), 3.11d
VIg	-	-	-	1.55s	3.99dt (697 and 8), 3.63 dt.
VIh	2.16 br.s	2.03m	1.66m	-	3.32d (698), 3.01d
VIIa	2.16 br.s	2.03m	1.65m	-	3.36d (696), 3.10d
VIIb	2.16 br.s	2.10m	1.67m	-	3.37d (695), 3.08d
VIIIa	2.16 br.s	2.09m	1.67m	-	3.35d (694), 3.07d
VIIIb	-	-	-	1.52s	3.94dt (692 and 8), 3.61 dt.

[B] STABILITY IN THE HUMAN SERUM

The compounds according to the present invention are slowly hydrolyzed to corresponding nucleoside 5'-hydrogenphosphonates in human blood serum.

- 5 The hydrolysis of the compounds according to the present invention was performed in human blood serum. The assay mixture containing 0.5 ml of 100 μM solution of the compound of the invention in formamide and 99.5 μl of 100% human fetal serum was incubated at 37°C. After certain intervals, the 5-μl samples were mixed with 45 μl of 10% trifluoroacetic acid, centrifuged for 10 min at 12,000 rpm, and the
- 10 supernatants were concentrated to 100 μl and analyzed by HPLC on a Silasorb C7 column (4 x 150 mm, 13 m) with a linear gradient of methanol from 0 to 80% in 0.05 M potassium phosphate buffer (pH 6.0) for 40 min. The flow rate was 0.5 ml/min. The extent of hydrolysis was assessed by measuring the amount of the product.

The results of the tests are shown in Table 4 hereinbelow.

TABLE 3

¹H NMR spectral data for "depot" groups in VI-VIII, 200 MHz; CD₃CN; δ, ppm; (J, Hz) and
³¹P NMR spectra of VI-VIII, 81 MHz, CD₃CN; δ, ppm, (J, Hz)

Comp.	<i>tert.</i> -CH (adam.), 3H	CH ₂ -C-O- P(adam.), 6H	CH ₂ (adam.) 6H	<i>tert.</i> -Bu, 9H	³¹ P NMR, 1P
VIa	2.16 br.s	2.06m	1.64m	-	3.36d (696), 3.08d
VIb	-	-	-	1.49s	3.99dt (697 and 8), 3.63 dt.
VIc	2.16 br.s	2.06m	1.63m	-	3.38d (694), 3.04d
VIId	2.16 br.s	2.08m	1.64m	-	3.33d (696), 3.07d
VIe	2.16 br.s	2.08m	1.64m	-	3.36d (694), 3.06d
VIIf	2.16 br.s	2.06m	1.63m	-	3.32d (695), 3.11d
VIg	-	-	-	1.55s	3.99dt (697 and 8), 3.63 dt.
VIh	2.16 br.s	2.03m	1.66m	-	3.32d (698), 3.01d
VIIa	2.16 br.s	2.03m	1.65m	-	3.36d (696), 3.10d
VIIb	2.16 br.s	2.10m	1.67m	-	3.37d (695), 3.08d
VIIIa	2.16 br.s	2.09m	1.67m	-	3.35d (694), 3.07d
VIIIb	-	-	-	1.52s	3.94dt (692 and 8), 3.61 dt.

[B] STABILITY IN THE HUMAN SERUM

The compounds according to the present invention are slowly hydrolyzed to corresponding nucleoside 5'-hydrogenphosphonates in human blood serum.

5 The hydrolysis of the compounds according to the present invention was performed in human blood serum. The assay mixture containing 0.5 ml of 100 μM solution of the compound of the invention in formamide and 99.5 μl of 100% human fetal serum was incubated at 37°C. After certain intervals, the 5-μl samples were mixed with 45 μl of 10% trifluoroacetic acid, centrifuged for 10 min at 12,000 rpm, and the
 10 supernatants were concentrated to 100 μl and analyzed by HPLC on a Silasorb C7 column (4 x 150 mm, 13 m) with a linear gradient of methanol from 0 to 80% in 0.05 M potassium phosphate buffer (pH 6.0) for 40 min. The flow rate was 0.5 ml/min. The extent of hydrolysis was assessed by measuring the amount of the product.

The results of the tests are shown in Table 4 hereinbelow.

15

THIS PAGE BLANK (USPTO)

TABLE 4

Hydrolysis of 50% of the compounds under the invention to corresponding nucleoside 5'-phosphonates

Compound	Time, min
VIa	60
VIb	5
VIc	55
VIId	35
VIe	42
VIIf	85
VIg	79
VIh	66
VIIa	150
VIIb	18
VIIIa	75
VIIIb	20

As seen in Table 4, in most cases the half-lives of the compounds according to the present invention are rather high.

TABLE 5

Concentration of the compounds under the invention inhibiting the HIV reproduction

Compounds of the invention	Concentration, inhibiting the virus reproduction by 50%, nM	concentration inhibiting the virus reproduction by 95%, nM	Concentration inhibiting cell growth for 50%, μ M
VIa	12	50	2800
VIb	45	210	3100
VIc	35	180	6800
VIId	5	65	120
VIe	0.8	6	2400
VIIf	14	95	5450
VIg	32	430	4300
VIh	88	590	4460
VIIa	88	460	3600
VIIb	56	235	2900
VIIc	92	430	280
VIIIa	33	160	890
VIIIb	37	165	165
AZT	3	16	1800
ddT	2500		>10000
ddC	2	14	80
FLT	2	8	1400
ddA	150	720	3000
ddG	400	1260	2480
d4T	80	310	2400
d4C	120	465	120
3TC	65	140	860

AZT - 3'-azido-2',3'-dideoxythymidine, FLT - 3'-fluoro-2',3'-dideoxythymidine

The data presented in Table 5 demonstrate that antiviral activity of the compounds according to the present invention is comparable to that of AZT and other nucleosides and is sometimes higher.

IC] ANTI-HIV ACTIVITY OF THE INVENTED COMPOUNDS

The data presented in Table 5 demonstrate antiviral activity and toxicity in HIV-infected MT4 cell culture of the compounds according to the present invention.

Antiviral activity and toxicity were studied according to [4,6].

TABLE 5

Concentration of the compounds under the invention inhibiting the HIV reproduction

Compounds of the invention	Concentration, inhibiting the virus reproduction by 50%, nM	concentration inhibiting the virus reproduction by 95%, nM	Concentration inhibiting cell growth for 50%, μ M
VIa	12	50	2800
VIb	45	210	3100
VIc	35	180	6800
VI d	5	65	120
VIe	0.8	6	2400
VI f	14	95	5450
VI g	32	430	4300
VI h	88	590	4460
VIIa	88	460	3600
VIIb	56	235	2900
VIIc	92	430	280
VIIIa	33	160	890
VIIIb	37	165	165
AZT	3	16	1800
ddT	2500		>10000
ddC	2	14	80
FLT	2	8	1400
ddA	150	720	3000
ddG	400	1260	2480
d4T	80	310	2400
d4C	120	465	120
3TC	65	140	860

AZT - 3'-azido-2',3'-dideoxythymidine, FLT - 3'-fluoro-2',3'-dideoxythymidine

The data presented in Table 5 demonstrate that antiviral activity of the compounds according to the present invention is comparable to that of AZT and other nucleosides and is sometimes higher.

[C] ANTI-HIV ACTIVITY OF THE INVENTED COMPOUNDS

The data presented in Table 5 demonstrate antiviral activity and toxicity in HIV-infected MT4 cell culture of the compounds according to the present invention.

Antiviral activity and toxicity were studied according to [4,6].

THIS PAGE BLANK (USPTO)

REFERENCES

The following publications, patents and patent applications referred to herein are incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety:

1. Tarusova N.B., Khorlin A.A., Krayevsky A.A., Korneyeva M.N., Nosik D.N., Kruglov I.V., Galegov G.A., Beabealashvili R.Sh., Inhibition of human immunodeficiency virus in cell culture by 3'-azido-2',3'-dideoxynucleosides, *Il.Biol. Russian*, 1989, 23, N6, 1716-1724.
2. Tarusova N.B., Kukhanova M.K., Krayevsky A.A., Karamov E.V., Lukashov V.V., Kornilayeva G.V., Rodina M.A., Galegov G.A., Inhibition of human immunodeficiency virus (HIV) production by 5'-hydrogenphosphonates of 3'-azido-2',3'-deoxynucleosides, *Nucleosides & Nucleotides*, 1991, 10, N1-3, 351-354.
3. Khorlin A.A., Tarusova N.B., Dyatkina N.B., Krayevsky A.A., Beabealashvili R.Sh., Galegov G.A., Zhdanov V.M., Korneyeva M.S., Nosik D.N., Maiorova S.N., Shobukhov V.M., Patent RF 1 548 182, 16.02.1992, US Patent 5,043,437, 27.8.1991; European Patent 0-354-246, 16.03.1994; Japan Patent N 0-354-246 B1, 08.25.1995; Corean Patent 106,957, 12.01.96.
4. Atrazheva E.D., Lukin M.A., Jasko M.V., Shushkova T.V., Tarusova N.B., Krayevsky A.A., Balzarini J., De Clercq E., 2',3'-O-Cyclic derivatives of ribonucleosides and their 5'-phosphonates: synthesis and anti-HIV activity, *Med.Chem.Res.*, 1991, 1, 155-165.
5. Karamov E.V., Lukashov V.V., Gorbachova A.P., Kornilayeva G.V., Tarusova N.B., Krayevsky A.A., Inhibition of HIV reproduction in cell cultures by 2',3'-dideoxynucleoside 5'-phosphites, *Il.Biol. Russian*, 1992, 26, N1, 201-207).

6. Krayevsky A.A., Tarusova N.B., Zhu Q.-Y., Vidal P., Chou T.-C., Baron P., Polsky B., Jiang X.-Y., Matulic-Adamic J., Rosenberg I., Watanabe K.A., 5'-Hydrogenphosphonates and 5'-methylphosphonates of sugar modified pyrimidine nucleosides as potential anti-HIV agents. *Nucleosides & Nucleotides*, 1992, 11, N2-4, 177-196.

7. Mc Guigan C., Bellevergue P., Jones B.C.N.M., Mahmood N., Hay A.J., Petrik J., Karpas A., Alkyl hydrogen phosphonate phosphonate derivatives of anti-HIV agent AZT may be less toxic than parent nucleosides, *Antiviral Chem., Chemother.*, 1994, 5, N4, 271-277.

8. Cardona V.M.F., Ayi A.L., Aubertin A.M., Guedj R., Anti-HIV activity of new compounds: Prodrug of D4T, *Antiviral Res.*, 1998, 37, N3 A52.

6. Krayevsky A.A., Tarusova N.B., Zhu Q.-Y., Vidal P., Chou T.-C., Baron P., Polsky B., Jiang X.-Y., Matulic-Adamic J., Rosenberg I., Watanabe K.A., 5'-Hydrogenphosphonates and 5'-methylphosphonates of sugar modified pyrimidine nucleosides as potential anti-HIV agents. *Nucleosides & Nucleotides*, 1992, 11, N2-4, 177-196.


7. Mc Guigan C., Bellevergue P., Jones B.C.N.M., Mahmood N., Hay A.J., Petrik J., Karpas A., Alkyl hydrogen phosphonate phosphonate derivatives of anti-HIV agent AZT may be less toxic than parent nucleosides, *Antiviral Chem., Chemother.*, 1994, 5, N4, 271-277.

8. Cardona V.M.F., Ayi A.L., Aubertin A.M., Guedj R., Anti-HIV activity of new compounds: Prodrug of D4T, *Antiviral Res.*, 1998, 37, N3 A52.


THIS PAGE BLANK (USPTO)

What is claimed is:


1. Adamantyl esters of 5'-O-hydrogenphosphonates of 3'-azido-2',3'-dideoxythymidine (phoshazide), 2',3'-dideoxythymidine, 3'-amino-2',3'-dideoxythymidine, 3'-fluoro-2',3'-dideoxythymidine, 2',3'-dideoxy-2',3'-didehydrothymidine, 2',3'-dideoxycytidine, 2',3'-dideoxyadenosine, 2',3'-dideoxyguanosine, L-3'-thiocytidine.
2. *tert*-Butyl esters of 5'-O-hydrogenphosphonates of 3'-azido-2',3'-dideoxythymidine (phoshazide), 2',3'-dideoxythymidine, 3'-amino-2',3'-dideoxythymidine, 3'-fluoro-2',3'-dideoxythymidine, 2',3'-dideoxy-2',3'-didehydrothymidine, 2',3'-dideoxycytidine, 2',3'-dideoxyadenosine, 2',3'-dideoxyguanosine, L-3'-thiocytidine.
3. Bicyclo[2,2,1] heptyl esters of 5'-O-hydrogenphosphonates of 3'-azido-2',3'-dideoxythymidine (phoshazide), 2',3'-dideoxythymidine, 3'-amino-2',3'-dideoxythymidine, 3'-fluoro-2',3'-dideoxythymidine, 2',3'-dideoxy-2',3'-didehydrothymidine, 2',3'-dideoxycytidine, 2',3'-dideoxyadenosine, 2',3'-dideoxyguanosine, L-3'-thiocytidine..
4. Compounds of the formulae VI-VIII:



VI



VII



VIII

wherein:

Nucleosides of D and L-series;

B = thymine, adenine, guanine, cytosine, uracyl, 5-fluorouracyl or 5-ethyluracyl;

R = N₃, NH₂, F, H or F;

X = S or O;

R = adamantyl-1, bicyclo[6,5,1] heptyl-1 or *tert.*-butyl.


B = thymine, adenine, guanine, cytosine, uracyl, 5-fluorouracyl or 5-ethyluracyl;

R = N₃, NH₂, F, H or F;

X = S or O;

R = adamantyl-1, bicyclo[6,5,1] heptyl-1 or *tert.*-butyl.

THIS PAGE BLANK (USPTO)

**FIG. 1**

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 00/00914

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07H19/10 C07H19/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>MCGUIGAN, C. ET AL.: "Alkyl hydrogen phosphonate derivatives of the anti-HIV agent AZT may be less toxic than the parent nucleoside analogue" ANTIVIR. CHEM. & CHEMOTHER., vol. 5, no. 4, 1994, pages 271-7, XP002105798 cited in the application the whole document</p> <p style="text-align: center;">-/--</p>	1-4

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

4 December 2000

Date of mailing of the international search report

18/12/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bardili, W

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 00/00914

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07H19/10 C07H19/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>MCGUIGAN, C. ET AL.: "Alkyl hydrogen phosphonate derivatives of the anti-HIV agent AZT may be less toxic than the parent nucleoside analogue"</p> <p>ANTIVIR. CHEM. & CHEMOTHER., vol. 5, no. 4, 1994, pages 271-7, XP002105798</p> <p>cited in the application</p> <p>the whole document</p> <p style="text-align: center;">--- -/-</p>	1-4

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *S* document member of the same patent family

Date of the actual completion of the international search

4 December 2000

Date of mailing of the international search report

18/12/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Bardili, W

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 00/00914

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CARDONA, V.M.F. ET AL.: "Synthesis and anti-HIV activity of some novel arylphosphonate and H-phosphonate derivatives" ANTIVIR. RES., vol. 42, no. 3, 1999, pages 189-196, XP000944433 cited in the application the whole document	1-4
Y	BERGSTROM, D.E. ET AL.: "Periodic-catastrophic transport of an adenosine 5'-monophosphate adamantadine conjugate through a model liquid membrane" NUCLEOSIDES, NUCLEOTIDES, vol. 10, 1991, pages 685-688, XP000944399 the whole document	1-4
Y	STEC, W.J. ET AL.: "Application of oxathiaphospholane method for the synthesis of oligodeoxyribonucleotide 5'-O-conjugates" NUCLEOSIDES, NUCLEOTIDES, vol. 17, 1998, pages 1977-82, XP000944463 the whole document	1-4
Y	WO 96 10030 A (ISIS PHARMACEUTICALS INC ;COOK PHILLIP DAN (US); TENG KELLY (US)) 4 April 1996 (1996-04-04) page 23, line 26 - line 35; example 16	1-4
A	US 5 043 437 A (KHORLIN ALEXANDR A ET AL) 27 August 1991 (1991-08-27)	

Form PCT/SA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/CA 00/00914

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9610030 A	04-04-1996	US 5608046 A AU 3892395 A US 5998603 A	04-03-1997 19-04-1996 07-12-1999
US 5043437 A	27-08-1991	SU 1548182 A DE 3888530 D EP 0354246 A JP 6089019 B KR 9609930 B WO 8906238 A	07-03-1990 21-04-1994 14-02-1990 09-11-1994 25-07-1996 13-07-1989

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)